



# Is Self-Face Recognition Special? Evidence from Lateralized Upside Down Faces

Yuan Hang Li and Eran Zaidel

Department of Psychology, University of California at Los Angeles



CNS 2008 Poster # E106


Contact: yuanli@ucla.edu

## Introduction

- Hemispheric Specialization: The Right Hemisphere (RH) is specialized for face recognition (Bradshaw & Nettleton, 1983).
- The Face Inversion Effect (FIE): Upright Faces are recognized better than Inverted Faces. This is attributed to holistic processing of Upright Faces and feature-dependent processing of Inverted Faces (Farah et al., 1995).
- The Hemispheric FIE (HFIE): The FIE is greater in the normal RH than in the normal Left Hemisphere (LH) (Leehey et al., 1978).
- We examined 1) whether RH specialization for recognizing faces in general, and 2) whether greater FIE in the RH, are true for recognizing one's own face.

## Method

- Participant: 34 UCLA undergraduate students (15 female) completed this experiment for course credit.
- Stimuli and Procedure: the subject's own face or a familiar friend's face were presented tachistoscopically (150ms) to the left or right visual hemifield, in an upright or inverted position, using the program E-Prime.
- Task: subjects judged whether the stimulus was 'self' or 'not self' and responded with both index fingers for 'self' and both middle fingers for 'not self.'



Task: Is the face shown "self" or "not self?"

Sample Stimuli





Self Upright Familiar Upright



Self Inverted Familiar Inverted

## Results

- A 2 x 2 x 2 (Face Type: Self, Familiar) x (VF: LVF, RVF) x (Inversion: Inverted, Upright) ANOVA was carried out. Dependent variables were latency and sensitivity (signal detection) of responses.
- Faces shown to the RH were recognized faster ( $p=0.002$ ) and with greater sensitivity ( $p<0.001$ ) than faces shown to the LH.



## Results cont'd

- Inverted Faces were recognized more slowly ( $p<0.001$ ) and less accurately ( $p<0.001$ ) than Upright Faces.
- There was a significant interaction (Face Type x VF) in latency ( $p=0.011$ ), showing that self-faces were recognized equally fast by both VF, whereas familiar faces were recognized faster in the LVF than the RVF (Top figure).
- There was a significant interaction (Inversion x VF) in sensitivity ( $p=0.02$ ), showing equal sensitivity in both VFs for recognizing Upright Faces but greater sensitivity in the LVF than in the RVF for recognizing Inverted Faces (Bottom figure).
- The interaction (Inversion x VF x Face Type) was not significant for either dependent variable.

## Discussion

- Our data showed RH specialization for recognizing familiar faces but equal ability of both hemispheres to recognize self-faces. Others also have found equal self-face recognition capacity in the two hemispheres (Uddin et al., 2005).
- We found that the FIE was larger in the RVF than the LVF, contrary to previous reports (e.g., Leehey et al., 1978).

## Conclusion

- Self-face recognition does not show the standard RH advantage for general face recognition. We conclude that self-face recognition is special.
- This extends previous studies that showed that self-face recognition is special by showing self-faces are recognized selectively better in the Upper Visual Fields (Li & Zaidel, 2006, 2007).

## References

- Bradshaw, J.L. & Nettleton, N. (1983). *Human cerebral asymmetry*. Englewood Cliffs, NJ: Prentice hall.
- Farah, M.J., Tanaka, J.W., & Drain, H.M. (1995). What causes the face inversion effect? *Journal of Experimental Psychology*, 21(3), 628-634.
- Leehey, S., Carey, S., Diamond, R., & Cahn, A. (1978). Upright and inverted faces: the right hemisphere knows the difference. *Cortex*, 14(3), 411-419.
- Li, Y.H. & Zaidel, E. (2006). *A Different Mechanism for Self-Face Recognition?* Poster session at the Cognitive Neuroscience Conference, San Francisco, CA.
- Li, Y.H. & Zaidel, E. (2007). *Upper and Lower Visual Field Differences in Self-Face Recognition*. Poster session at the Cognitive Neuroscience Society Conference, New York, NY.
- Uddin, L.Q., Rayman, J., & Zaidel, E. (2005). Split-brain reveals separate but equal self-recognition in the two cerebral hemispheres. *Consciousness and Cognition*, 14(3), 633-640.